



**CANADIAN ACADEMY OF SPORT MEDICINE  
ACADÉMIE CANADIENNE DE MÉDECINE DU  
SPORT**

*“Committed to Excellence • L’excellence dans la pratique”*

## **DISCUSSION PAPER**

### ***Head Injuries and Concussions in Soccer***

J. Scott Delaney, MDCM, FRCPC, FACEP and  
Renata Frankovich BMath, MD, FCFP, Dip Sport Med

This discussion paper was updated and reviewed by members of the Canadian Academy of Sport Medicine (CASEM) Sport Safety Committee. This discussion paper was approved by the CASEM Board of Directors as a CASEM discussion paper in September, 2010.

#### **1.0 INTRODUCTION**

Soccer has not always been perceived as a high risk sport for head injuries or concussions.<sup>1,2,3,4,5</sup> However, recent research suggests that soccer players have head injury and concussion rates similar to football and ice hockey.<sup>5,6,7, 8, 9, 10, 11,12, 13,14,15,16,17,18,19</sup> [1] The Canadian Academy of Sport and Exercise Medicine (CASEM) has undertaken this discussion paper to review the literature and to provide recommendations to the public to decrease the risk of head injuries and concussions in the sport of soccer.

#### **2.0 DEFINITIONS**

**Head Injury** – a traumatic injury to the head which is usually evident on clinical examination. It is usually characterized by ecchymoses, hematoma, lacerations, deformities, or cerebral spinal fluid leakage.<sup>1</sup> A head injury may constitute a minor diagnosis, such as a scalp laceration, or it may comprise a more serious diagnosis such as a concussion or a skull fracture.

**Concussion** – a form of head injury characterized by any alteration in cerebral function and caused by a direct or indirect (rotation) force transmitted to the head. It results in one or more of the following acute signs or symptoms: a brief loss of consciousness, light-headedness, vertigo, cognitive and memory dysfunction, tinnitus, blurred vision, difficulty concentrating, amnesia, headache, nausea, vomiting, photophobia or a balance disturbance. Delayed signs and symptoms may also include sleep irregularities, fatigue, personality changes, inability to perform usual daily activities, depression or lethargy.<sup>2,3,4, [2] [3]</sup>

## **3.0 RECOMMENDATIONS**

### **3.1 Soccer should be regarded as a contact sport in which players are at risk for head injuries and concussions.**

Studies have shown that almost half of varsity level university soccer players may experience symptoms from a concussion during just one fall season,<sup>5</sup> and almost two thirds of these players may experience symptoms from a concussion over one full year of soccer participation.<sup>6</sup> When comparing different sports, research has shown that soccer has head injury and concussion rates similar to football and ice hockey.<sup>5,6,7,8, 9,10,11,12, 13, 14,15,16,17,18,19</sup> These injury rates are comparable, not only for elite athletes participating in soccer, football and ice hockey, but also for recreational and community athletes participating in these sports.<sup>20</sup>

### **3.2 Safe play and respect for one's opponent should be emphasized.**

Concussions during participation in soccer are caused most frequently by collisions and contact among players.<sup>8</sup>[4-7] The rules of soccer do not allow any player to engage in play which might endanger the safety of another player,<sup>21</sup>note: 21 is updated in references and referees should continue to ensure that reckless and potentially harmful actions to others are not permitted during the game[8, 9]. The spirit of fair play and respect for one's opponent should be emphasized by parents, coaches and referees.

### **3.3 Players, parents and coaches should be aware of the signs and symptoms of a concussion.**

Research has revealed that the majority of sport related concussions may go unrecognized.<sup>5,6, 22,23</sup> The need to identify an acute concussion and prevent a player from returning to competition is vital because this may prevent a more severe concussion or life threatening complications if a player receives another blow to the head while still suffering symptoms of a concussion.<sup>23, 24,25,26,27,28,29,,30,31,32</sup> It is also important to accurately diagnose a concussion because multiple concussions may result in longer and more severe episodes of functional disability.<sup>12,33</sup> [10] Repeated concussions may also result in progressive and cumulative neurologic and neuropsychologic impairment.<sup>16,34,35</sup> [11, 12]

In an effort to provide a safer environment for athletes, it is important that players, parents and coaches be reminded of the common signs and symptoms of a concussion (see definition above). The need to err on the side of caution when diagnosing a concussion should be emphasized (“When in doubt, sit them out!”).<sup>36</sup>, [3]

### **3.4 Any athlete suspected of having sustained a concussion should be removed from competition. The athlete should not return to play until they are examined by a medical doctor familiar with diagnosing and treating sport related concussions.**

When diagnosing and deciding disposition for a concussed athlete, concussion guidelines such as those by the Concussion in Sport Group, can help guide return to play decisions.<sup>2,36</sup> [2] [3] Parents, players, coaches and health care providers should be reminded of the critical points of the guidelines:

- a. Players who show any signs and symptoms of a concussion should not be allowed to return to play in the current game or practice.
- b. Concussed players should not be left alone during the immediate period following a concussion. Close monitoring for deterioration is essential in the initial post-concussion period and should continue at least until the player's symptoms have stabilized.
- c. The concussed athlete should be evaluated by a medical doctor with knowledge and experience in concussion management.
- d. Return to play must follow a gradual process. It must be monitored by a medical doctor with knowledge and experience in concussion management.

**1. 3.5 Only soccer balls which are the appropriate size, in good condition, and inflated appropriately should be used.**

Just as all parents and players would insist on proper sized protective equipment and athletic gear, it should be remembered that soccer balls also come in different sizes. Using different size balls is intended to prevent injury and make the balls more manageable for different size players[13]. The smaller balls are lighter and easier for children to manipulate. The smallest size (No. 3) is used for children under 10 years of age, the medium sized ball (No. 4) is used for juniors (10 to 14 years of age), and the largest ball (No. 5) is used for athletes over 14 years of age.<sup>18,37,38</sup>

Soccer balls should be checked frequently for proper inflation pressures. The pressure inside a ball may change over time because of frequent use or because of changes in temperature and weather conditions. The recommended inflation pressure is usually found on the ball. Hyperinflation of soccer balls has been proposed as a possible risk factor for headaches and possibly concussions in soccer players.<sup>7,39,40</sup>

Most modern soccer balls are made from synthetic materials which are water resistant.<sup>10</sup> Older balls made of leather or newer synthetic balls with cracks in the material should not be used in wet conditions as the weight of a wet ball may increase by 20% or more and increase the amount of force absorbed by the cranium during heading.<sup>18,41</sup>

**3.6 Children should minimize heading the ball until there is a better understanding of the immediate and possible long-term effects of heading and until they sufficiently master the proper heading techniques.**

Due to children's immature anatomy and inexperience, many believe that children may be more at risk for head and brain injuries than adults.<sup>19, 40,42</sup> Compared to adults, children have proportionately larger heads but thinner skulls to protect the brain.<sup>9</sup>

Younger soccer players also have weaker neck muscles which, when developed, may help to absorb and dissipate the forces applied to the head.<sup>2,3, 10, 16,18,43,44,45</sup> Children may also be less adept at the techniques of heading a ball, which may decrease the amount of energy transmitted to the brain.<sup>13,16,</sup>

Whether heading is a significant mechanism of injury leading to concussions in adults is less controversial now than it once was. Studies addressing mechanism of injuries showed that the ball was involved in less than 25% of all cases of concussions and few, if any, concussions occurred during uninterrupted purposeful heading.<sup>8, 10, 37</sup> [7] Other studies showed that neither the average number of headings per game, nor considering oneself to be someone who heads the ball frequently, was a significant risk factor for concussions.<sup>5,6</sup> While there is less fear that unimpeded heading the ball causes a significant number of concussions, research has shown that heading duels or aerial challenges between two players preparing to head the ball is a common scenario for head injury and concussions [5] [6]. The question of possible long-term effects of heading remains controversial. European studies showed neuropsychological and neuroanatomic changes in active and retired soccer players when compared to controls.<sup>12,14,16,17,</sup> North American studies tended not to show these deleterious effects.<sup>46,47,</sup> [14, 15].

Suggestions for dealing with heading include outlawing heading in young age groups, limiting its use in young age groups or continuing with no special precautions as in the past.<sup>10,39</sup> Until long-term risks of heading soccer balls are more clearly elucidated, it is prudent to limit the amount of heading that young children perform, at least until an age when they begin to master the heading techniques and their neck and skull anatomy are more mature.<sup>42</sup>

### **3.7 Proper heading techniques should be taught in a supervised and controlled setting.**

It is well accepted that when proper methods of heading a soccer ball are used, these techniques may decrease the amount of force transmitted to the brain.<sup>10,17,</sup> These techniques should be taught by a qualified individual with good knowledge of different methods of heading a soccer ball in order to lessen the force applied to the cranium during heading.<sup>13,39,42</sup> This is especially important when children first learn to head the soccer ball. Some experts suggest using a specially designed lightweight ball or small ball suspended from a rope (“piñata model”) to initially illustrate proper body positioning and contact points.<sup>38</sup> Neck musculature strengthening should be emphasized because many believe that strong neck muscles may help dissipate the energy applied to the head after contact with the soccer ball.<sup>6,10,15,40,41,</sup> although this has not yet been proven in a prospective study. [16]

### **3.8 Goalposts should be appropriately padded and anchored to the ground.**

Incidents with goalposts have resulted in many serious head injuries and even several deaths. In the past few decades, most fatalities from soccer-related injuries have come from traumatic contact with goalposts.<sup>42,48,49</sup> Portable goalposts need to be anchored in a secure and approved fashion as death or serious head and neck injury can easily occur

from goalposts that fall forward onto an unsuspecting child or adolescent.<sup>50</sup> In the event of contact with goalposts, portable and permanently installed posts should have adequate padding to reduce injuries. Data has shown that padding goalposts can decrease the number of injuries from contact with goalposts.<sup>51</sup>

While different shapes are permitted for goalposts, further research is required to validate the belief that round or elliptical shaped goalposts are safer than square or rectangular shaped goalposts.

### **3.9 Goalkeeper is the position most at risk for concussion and these players should be protected accordingly.**

Numerous studies have shown that goalkeepers are the players most at risk for concussion.<sup>5, 6, 52, 53, 54</sup> The role of a goalkeeper is unique. They are expected to stop balls, often kicked from a short distance. These balls can travel at speeds in excess of 120 km/hr.<sup>16</sup> They are also vulnerable to being kicked, kneed or elbowed in the head as players converge on the goal, or as they cover up a ball at or near ground level. Players, coaches and referees should be aware of the vulnerable situations in which goalies are placed. Respect for their safety should be emphasized and the rules that protect them should be strictly enforced at all times. Due to their increased risk of concussions, goalies may be the position most likely to benefit from a proven and effective form of headgear.

### **3.10 Mouth guards should be worn during participation in soccer.**

Current evidence that mouth guards prevent sport related concussions is poor. While a few studies suggested a possible benefit,<sup>55, 56, 57, 58</sup> others failed to show any benefit.<sup>6, 59, 60</sup> While the effectiveness of mouth guards in preventing concussions is controversial, the effectiveness in preventing orofacial and mandibular injuries is not.<sup>59, 61, 62, 63</sup> Wearing a mouth guard is a proven and reliable way to prevent orofacial and mandibular injuries. The proper use of well-made mouth guards should be recommended for its proven role in preventing orofacial and mandibular injuries and for its possible, if yet unproven, benefit of preventing concussions.

### **3.11 There may be a role for the use of protective headgear in soccer for those athletes seeking more protection from concussion and head injuries.**

The use of protective headgear has been suggested by some as a possible means of decreasing the number of head injuries and concussions in soccer.<sup>9, 54</sup> Children, goalies and players with a past history of head injuries or concussions have been suggested as the groups most likely to benefit from its use.<sup>9</sup> Preliminary laboratory and field research has shown its use may decrease the number of concussions and soft tissue injuries on the head. [17] [18] The international governing body of soccer now permits the use of these products,<sup>21</sup> ( 21-this ref is updated) and several leagues and soccer facilities in North America have already mandated the use of protective headgear for soccer.<sup>47, 64</sup> The American Society for Testing and Materials (ASTM) has created a national safety

standard and certification process for protective headgear for soccer in the United States. This standard addresses the safety and effectiveness of the products being sold. This recognized national safety certification standard allows the public to know that any certified product has met or exceeded an acceptable level of quality and safety for soccer. Further field studies are necessary to better define the extent of protective effects of headgear for soccer.

## **4.0 CONCLUSION**

The need to inform players, parents and coaches about fair and safe play is vital. The recommendations above are intended to decrease head injuries and concussions during participation in soccer, but recommendations are useless without education. Along with further research to improve understanding of sport related head injuries and concussions, education may be as valuable as any piece of protective equipment in preventing these injuries.

## 5.0 REFERENCES

1. Gottesfeld SH, Jagoda A. Mild head trauma: appropriate diagnosis and management. *Emergency Medicine Practice* 2000;2:1-24.
2. Aubry M, Cantu R, Dvorak J, et al. Summary and agreement statement of the 1st international symposium on concussion in sport, vienna 2001. *Clin J Sport Med* 2002; 12:6-11.
3. Johnston KM, McCrory P, Mohtadi NG, et al. Evidence-Based review of sport-related concussion: clinical science. *Clin J Sport Med* 2001;11:150-9.
4. Wojtys EM, Hovda D, Landry G, et al. Current concepts. Concussion in sports. *Am J Sports Med* 1999;27:676-87.
5. Delaney JS, Lacroix VJ, Gagne C, et al. Concussions among university football and soccer players: a pilot study. *Clin J Sport Med* 2001;11:234-40.
6. Delaney JS, Lacroix VJ, Leclerc S, et al. Concussions among university football and soccer players. *Clin J Sport Med* 2002;12:331-8.
7. Barnes BC, Cooper L, Kirkendall DT, et al. Concussion history in elite male and female soccer players. *Am J Sports Med* 1998;26:433-8.
8. Boden BP, Kirkendall DT, Garrett WE, Jr. Concussion incidence in elite college soccer players. *Am J Sports Med* 1998;26:238-41.
9. Delaney JS, Drummond R. Has the time come for protective headgear for soccer? *Clin J Sport Med* 1999;9:121-3.
10. Kirkendall DT, Jordan SE, Garrett WE. Heading and head injuries in soccer. *Sports Med* 2001;31:369-86.
11. Matser JT, Kessels AG, Jordan BD, et al., Chronic traumatic brain injury in professional soccer players. *Neurology* 1998;51:791-6.
12. Matser EJ, Kessels AG, Lezak MD, et al. Neuropsychological impairment in amateur soccer players. *Jama* 1999;282:971-3.
13. Schmidt-Olsen S, Jorgensen U, Kaalund S, et al. Injuries among young soccer players. *Am J Sports Med* 1991;19:273-5.
14. Sortland O, Tysvaer AT. Brain damage in former association football players. An evaluation by cerebral computed tomography. *Neuroradiology* 1989;31:44-8.

15. Tysvaer A, Storli O. Association football injuries to the brain. A preliminary report. *Br J Sports Med* 1981;15:163-6.
16. Tysvaer AT, Storli OV. Soccer injuries to the brain. A neurologic and electroencephalographic study of active football players. *Am J Sports Med* 1989;17:573-8.
17. Tysvaer AT, Lochen EA. Soccer injuries to the brain. A neuropsychologic study of former soccer players. *Am J Sports Med* 1991;19:56-60.
18. Tysvaer AT, Head and neck injuries in soccer. Impact of minor trauma. *Sports Med* 1992;14:200-13.
19. Pediatrics TAAo. A policy statement. *The Physician and Sportsmedicine*.1988;16:165-9.
20. Delaney JS. Head injuries presenting to Emergency Departments in the United States from 1990 to 1999 for ice hockey, soccer and football. *Clin J Sport Med* 2004;14:80-7.
21.  
[http://www.fifa.com/mm/document/affederation/generic/81/42/36/laws\\_of\\_the\\_game\\_2010\\_11\\_e.pdf](http://www.fifa.com/mm/document/affederation/generic/81/42/36/laws_of_the_game_2010_11_e.pdf)
22. Delaney JS, Lacroix VJ, Leclerc S, et al. Concussions during the 1997 Canadian Football League season. *Clin J Sport Med* 2000;10:9-14.
23. Leclerc S, Lasseonde M, Delaney JS, et al. Recommendations for grading of concussion in athletes. *Sports Med* 2001;31:629-36.
24. Cantu RC. Return to play guidelines after a head injury. *Clin Sports Med* 1998;17:45-60.
25. Clarke KS. Epidemiology of athletic head injury. *Clin Sports Med*, 1998;17:1-12.
26. Cantu RC Cerebral concussion in sport. Management and prevention. *Sports Med* 1992;14:64-74.
27. McCrory PR, Berkovic SF. Second impact syndrome. *Neurology* 1998;50:677-83.
28. Torg JS. Epidemiology, pathomechanics, and prevention of athletic injuries to the cervical spine. *Med Sci Sports Exerc* 1985;17:295-303.
29. Cantu RC. Guidelines for return to contact sports after a cerebral concussion. *Phys Sportsmed* 1986;14:75-83.

30. Ommaya AK, Gennarelli TA. Cerebral concussion and traumatic unconsciousness. Correlation of experimental and clinical observations of blunt head injuries. *Brain* 1974; 97:633-54.
31. Roberts W. Who plays? Who sits? Managing concussion on the sidelines. *Phys Sportsmed* 1992;20:66-72.
32. Wilberger JE, Jr, Maroon JC. Head injuries in athletes. *Clin J Sports Med* 1989;8:1-9.
33. Lindsay KW, McLatchie G, Jennett B. Serious head injury in sport. *Br Med J* 1980;281:789-91.
34. Warren WL, Jr, Bailes JE. On the field evaluation of athletic neck injury. *Clin Sports Med* 1998;17:99-110.
35. Kelly JP Rosenberg JH. Diagnosis and management of concussion in sports. *Neurology*, 1997;48:575-80.
36. Medicine CAoS, Guidelines for assessment and management of sport-related concussion. Canadian Academy of Sport Medicine Concussion Committee. *Clin J Sport Med*, 2000;10:209-11.
37. Smolaka VN. Medical aspects of heading the ball. *Phys Sportsmed* 1984;12:127-131.
38. Personal Communication - Gittens R. Teaching heading to young soccer players. Editor. 2003: Ottawa.
39. Green GA, Jordan SE. Are brain injuries a significant problem in soccer? *Clin J Sports Med* 1998;17:795-809,viii.
40. Queen RM, Weinhold PS et al. Theoretical study of the effect of ball properties on impact force in soccer heading. *Med Sci Sports Exerc* 2003;35: 2069-76.
41. Smolaka VJ. Medical aspects of heading the ball in soccer. *Phys Sportsmed* 1984; 12:127-31.
42. Pediatrics AAo, Injuries in youth soccer: a subject review. American Academy of Pediatrics. Committee on Sports Medicine and Fitness. *Pediatrics*, 2000;105:659-61.
43. Cantu RC. Head injuries in sport. *Br J Sports Med* 1996;30:289-96.
44. Cantu RC. Athletic head injuries. *Clin J Sports Med* 1997;16:531-42.

45. McCrory P, Johnston KM, Mohtadi NG, et al. Evidence-based review of sport-related concussion: basic science. *Clin J Sport Med* 2001;11:160-5.
46. Echemendia RJ, Putukian M, Mackin RS, et al. Neuropsychological test performance prior to and following sports-related mild traumatic brain injury. *Clin J Sport Med* 2001; 11:23-31.
47. Guskiewicz KM, Marshall SW, Broglio SP, et al. No evidence of impaired neurocognitive performance in collegiate soccer players. *Am J Sports Med* 2002;30:157-62.
48. Blond L Hansen LB. Injuries caused by falling soccer goalposts in Denmark. *Br J Sports Med* 1999;33:110-2.
49. Commission USCPS, United States Consumer Product Safety Commission Injuries associated with soccer goalposts: United States, 1979-1993. *Morb Mortal Wkly Rep* 1994;43:153-5.
50. Injuries associated with soccer goalposts-United States, 1979-1993. *Morb Mortal Wkly Rep* 1994;43:153-5.
51. Janda DH, Bir C, Wild B, et al. Goal post injuries in soccer. A laboratory and field testing analysis of a preventive intervention. *Am J Sports Med* 1995;23:340-4.
52. Tucker AM. Common soccer injuries. Diagnosis, treatment and rehabilitation. *Sports Med* 1997;23:21-32.
53. Sullivan JA, Gross RH, Grana WA, et al. Evaluation of injuries in youth soccer. *Am J Sports Med* 1980;8:325-7.
54. Dailey SW, Barsan WG. Head injuries in soccer. A case for protective headgear? *Phys Sportmed* 1992;20:79-85.
55. Barth JT, Freeman JR, Winters JE. Management of sports-related concussions. *Dent Clin North Am* 2000;44:67-83.
56. Chapman PJ. Orofacial injuries and mouth guards: a study of the 1984 Wallabies. *Br J Sports Med* 1985;19:93-5.
57. Hickey JC, Morris AL, Carlson LD, et al. The relation of mouth protectors to cranial pressure and deformation. *J Am Dent Assoc* 1967;74:735-40.
58. Stenger JM, Lawton EA, Wright JM, et al. Mouth guards: protection against shock to head, neck and teeth. *Basal Facts* 1987;9:133-9.

59. Labella CR, Smith BW, Sigurdsson A. Effect of mouth guards on dental injuries and concussions in college basketball. *Med Sci Sports Exerc* 2002;34:41-4.
60. McCrory P. Do mouth guards prevent concussion? *Br J Sports Med* 2001;35:81-2.
61. Chalmers DJ. Mouth guards. Protection for the mouth in rugby union. *Sports Med* 1998;25:339-49.
62. Ranalli DN, Demas PN. Orofacial injuries from sport: preventive measures for sports medicine. *Sports Med* 2002;32:409-18.
63. Newsome PR, Tran DC, Cooke MS. The role of the mouthguard in the prevention of sports-related dental injuries: a review. *Int J Paediatr Dent* 2001;11:396-404.
64. [http://www.astm.org/SNEWS/AUGUST\\_2000/aug\\_soccer.html](http://www.astm.org/SNEWS/AUGUST_2000/aug_soccer.html). 2000.

### NEW References

1. Straume-Naesheim, T.M., et al., *Do minor head impacts in soccer cause concussive injury? A prospective case-control study*. *Neurosurgery*, 2009. **64**(4): p. 719-25; discussion 725.
2. McCrory, P., et al., *Summary and Agreement Statement of the 2nd International Conference on Concussion in Sport, Prague 2004*. *Clin J Sport Med*, 2005. **15**(2): p. 48-55.
3. McCrory, P., et al., *Consensus statement on Concussion in Sport 3rd International Conference on Concussion in Sport held in Zurich, November 2008*. *Clin J Sport Med*, 2009. **19**(3): p. 185-200.
4. Delaney, J.S., V. Puni, and F. Rouah, *Mechanisms of injury for concussions in university football, ice hockey, and soccer- a pilot study*. *Clin J Sport Med*, 2006. **16**(2): p. 162-5.
5. Fuller, C.W., A. Junge, and J. Dvorak, *A six year prospective study of the incidence and causes of head and neck injuries in international football*. *Br J Sports Med*, 2005. **39 Suppl 1**: p. i3-9.
6. Andersen, T.E., et al., *Mechanisms of head injuries in elite football*. *Br J Sports Med*, 2004. **38**(6): p. 690-6.
7. Withnall, C., et al., *Biomechanical investigation of head impacts in football*. *Br J Sports Med*, 2005. **39 Suppl 1**: p. i49-57.
8. Fuller, C.W., A. Junge, and J. Dvorak, *An assessment of football referees' decisions in incidents leading to player injuries*. *Am J Sports Med*, 2004. **32**(1 Suppl): p. 17S-22S.
9. Fuller, C.W., et al., *An assessment of player error as an injury causation factor in international football*. *Am J Sports Med*, 2004. **32**(1 Suppl): p. 28S-35S.
10. De Beaumont, L., et al., *Long-term and cumulative effects of sports concussion on motor cortex inhibition*. *Neurosurgery*, 2007. **61**(2): p. 329-36; discussion 336-7.

11. De Beaumont, L., et al., *Long-term electrophysiological changes in athletes with a history of multiple concussions*. *Brain Inj*, 2007. **21**(6): p. 631-44.
12. De Beaumont, L., et al., *Brain function decline in healthy retired athletes who sustained their last sports concussion in early adulthood*. *Brain*, 2009. **132**(Pt 3): p. 695-708.
13. Shewchenko, N., et al., *Heading in football. Part 3: effect of ball properties on head response*. *Br J Sports Med*, 2005. **39 Suppl 1**: p. i33-9.
14. Putukian, M., R.J. Echemendia, and S. Mackin, *The acute neuropsychological effects of heading in soccer: a pilot study*. *Clin J Sport Med*, 2000. **10**(2): p. 104-9.
15. Schmitt, D.M., et al., *Effect of an acute bout of soccer heading on postural control and self-reported concussion symptoms*. *Int J Sports Med*, 2004. **25**(5): p. 326-31.
16. Mansell, J., et al., *Resistance training and head-neck segment dynamic stabilization in male and female collegiate soccer players*. *J Athl Train*, 2005. **40**: p. 310-9.
17. Withnall, C., et al., *Effectiveness of headgear in football*. *Br J Sports Med*, 2005. **39 Suppl 1**: p. i40-8; discussion i48.
18. Delaney, J.S., et al., *The Effect of Protective Headgear on Head Injuries and Concussions in Adolescent Football (Soccer) Players*. *Br J Sports Med*, 2007.